

infection, transfection, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding sequence combination that is the same as, or different from a control sequence/DNA coding sequence combination found in the native cell.

5 **Vector**

As used herein, the term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression  
10 vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

**Expression Cassette**

Accordingly, an "expression cassette" or "expression vector" is a nucleic acid construct generated recombinantly or synthetically, with a series  
15 of specified nucleic acid elements that permit transcription of a particular nucleic acid in a target cell. The recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA, virus, or nucleic acid fragment. Typically, the recombinant expression cassette portion of an expression vector includes, among other sequences, a  
20 nucleic acid sequence to be transcribed and a promoter.

**Plasmid**

As used herein, the term "plasmid" refers to a circular double-stranded (ds) DNA construct used as a cloning vector, and which forms an extrachromosomal self-replicating genetic element in many bacteria and  
25 some eukaryotes.

As used herein, the term "selectable marker-encoding nucleotide sequence" refers to a nucleotide sequence which is capable of expression in mammalian cells and where expression of the selectable marker confers to cells containing the expressed gene the ability to grow in the presence of a  
30 corresponding selective agent.

### Promoter

As used herein, the term "promoter" refers to a nucleic acid sequence that functions to direct transcription of a downstream gene. The promoter will generally be appropriate to the host cell in which the target gene is being expressed. The promoter together with other transcriptional and translational regulatory nucleic acid sequences (also termed "control sequences") are necessary to express a given gene. In general, the transcriptional and translational regulatory sequences include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences.

"Chimeric gene" or "heterologous nucleic acid construct", as defined herein refers to a non-native gene (*i.e.*, one that has been introduced into a host) that may be composed of parts of different genes, including regulatory elements. A chimeric gene construct for transformation of a host cell is typically composed of a transcriptional regulatory region (promoter) operably linked to a heterologous protein coding sequence, or, in a selectable marker chimeric gene, to a selectable marker gene encoding a protein conferring antibiotic resistance to transformed cells. A typical chimeric gene of the present invention, for transformation into a host cell, includes a transcriptional regulatory region that is constitutive or inducible, a signal peptide coding sequence, a protein coding sequence, and a terminator sequence. A chimeric gene construct may also include a second DNA sequence encoding a signal peptide if secretion of the target protein is desired.

A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA encoding a secretory leader, *i.e.*, a signal peptide, is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate

translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites.

- 5 If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, that may or may not include regions preceding and following the coding region, e.g. 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

The gene may encode therapeutically significant proteins or peptides, such as growth factors, cytokines, ligands, receptors and inhibitors, as well as vaccines and antibodies. The gene may encode commercially important industrial proteins or peptides, such as enzymes, e.g., proteases, carbohydrases such as amylases and glucoamylases, cellulases, oxidases and lipases. The gene of interest may be a naturally occurring gene, a mutated gene or a synthetic gene.

A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under moderate to high stringency hybridization and wash conditions. Hybridization conditions are based on the melting temperature ( $T_m$ ) of the nucleic acid binding complex or probe. For example, "maximum stringency" typically occurs at about  $T_m - 5^\circ\text{C}$  ( $5^\circ$  below the  $T_m$  of the probe); "high stringency" at about  $5 - 10^\circ$  below the  $T_m$ ; "intermediate stringency" at about  $10 - 20^\circ$  below the  $T_m$  of the probe; and "low stringency" at about  $20 - 25^\circ$  below the  $T_m$ . Functionally, maximum stringency conditions may be used to identify sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to